

Ladder operators for the Coulomb potential

J López-Bonilla

The International Institute of Integral Human Sciences, 1974 de Maisonneuve West, Montréal, Qué, Canada H3H 1K5

V Gaftoi and J J Peña

Area de Física, División de CBI, Universidad Autónoma Metropolitana-Azcapotzalco Apdo, Postal 16-306, 02200 México, D.F.

e-mail address : gboz@hp9000a1.uam.mx

Received 13 July 1999, accepted 16 November 1999

Abstract : The ladder operators for the non-relativistic quantum Coulomb problem are obtained by solving a linear system of first-order differential equations. This approach being simpler than standard ones.

Keywords : Alternative approach to factorization method; ladder operators, hydrogenlike wave functions.

PACS Nos. : 03.65.Fd, 02.90.+p

The Schrödinger equation for a hydrogen-like atom with nuclear charge Ze is [1] ($h = m = 1$) :

$$-\frac{1}{2} \left[\frac{d^2}{dr^2} - \frac{l(l+1)}{r^2} \right] \psi_{nl}(r) - \frac{Ze^2}{r} \psi_{nl}(r) = \frac{Z^2 e^4}{2n^2} \psi_{nl}(r), \quad (1)$$

where Z is the atomic number, e the electron charge, $\frac{1}{r} \psi_{nl}$ is the hydrogenic radial wave function, $n = 1, 2, 3, \dots$ is the principal quantum number, and $l = 0, 1, \dots, n-1$ is the orbital quantum number.

This paper has the goal of determining ladder operators A_{nl}^{\pm} for a given value of n , such that

$$\psi_{n,l\pm 1}(r) = A_{nl}^{\pm} \psi_{nl}(r). \quad (2)$$

In this way, given the state ψ_{n0} we can obtain all the states ψ_{nl} applying the operator A_{nl}^+ an appropriate number of times; for example, $\psi_{n1} = A_{n0}^+ \psi_{n0}$ and so on. Otherwise, we might use A_{nl}^- to get all the ψ_{nl} starting with $\psi_{n,n-1}$.

There are various methods proposed for getting this kind of ladder operators; for example, the procedure used in [2] requires the explicit expression of ψ_{nl} , but this is not necessary in the technique proposed in this work. It neither

needs to invoke the factorization method [3–9] to deduce A_{nl}^{\pm} as is done by certain authors [3,10]. There are various other methods [11] where the ladder operators are obtained taking advantage of the mapping [12–16] relating the Coulomb with the Morse [7,17–22] problem. In this work we take a different approach and propose that the ladder operators may be written as

$$A_{nl}^+ = f(r) \frac{d^2}{dr^2} + g(r) \frac{d}{dr} + h(r) \quad (3)$$

and show that eq. (1) and (2) lead to a system of linear first-order differential equations for f , g and h . As a matter of fact, we explicitly do the calculations only for A_{nl}^+ since those for A_{nl}^- follow a similar process.

If we define the operator

$$B_{nl} = -\frac{1}{2} \left[\frac{d^2}{dr^2} - \frac{l(l+1)}{r^2} \right] - \frac{Ze^2}{r} - \frac{Z^2 e^4}{2n^2} \quad (4)$$

then eq. (1) can be rewritten as

$$B_{nl} \psi_{nl} = 0. \quad (5)$$

Use of eq. (5) for $\psi_{n,l+1}$ together with the relevant component of eq. (2) allows as to obtain $B_{n,l+1} A_{nl}^+ \psi_{nl} = 0$, that is

$$\{ [B_{n,l+1}, A_{nl}^+] + A_{nl}^+ B_{n,l+1} \} \psi_{nl} = 0, \quad (6)$$

where $[C,D]$ stands for the commutator of C and D . From eq. (4) it follows that $B_{n,l+1} = B_{nl} + (l+1)/r^2$, which may be substituted into (6) and together with eq. (5) it leads to

$$\vartheta_{nl}\psi_{nl} = 0,$$

$$\text{with } \vartheta_{nl} = [B_{n,l+1}, A_{nl}^+] + (l+1)A_{nl}^+ \frac{1}{r^2}. \quad (7)$$

Eq. (7) is satisfied if we impose the condition $\vartheta_{nl} = 0$, which in union with relations (3) and (5) leads to the following set of equations

$$f' = 0, \quad (8)$$

$$g' + \frac{1}{2}f'' - (l+1)\frac{f}{r^2} = 0, \quad (9)$$

$$h' + \frac{2}{r^2}\left[Ze^2 + \frac{l(l+1)}{r}\right]f + \frac{1}{2}g'' - (l+1)\frac{g}{r^2} = 0, \quad (10)$$

$$\frac{1}{5}h'' - (l+1)\frac{h}{r^2} - \frac{Ze^2}{r^2}\left(\frac{2f}{r} - g\right) + \frac{l(l+1)}{r^3}\left(\frac{3f}{r} - g\right) = 0, \quad (11)$$

where the primes represent r -derivatives. Eq. (8) imply that $f(r) = C_1$, a constant, which together with (9) implies a first order differential equation for $g(r)$ whose solution is $g(r) = C_2 - (l+1)C_1/r$. The last function $h(r)$ in eq. (3) can now be obtained by solving (10):

$$h(r) = C_3 + \frac{1}{r}[2Ze^2C_1 - C_2(l+1)] + \frac{3C_1}{2} \frac{l(l+1)}{r}.$$

Finally, the expressions for f , g and h are substituted into equation (11) to get $C_1 = 0$ and $C_2 = (l+1)C_3/Ze^2$, from which we deduce

$$A_{nl}^+ = \frac{(l+1)}{Ze^2}C_3\left(\frac{d}{dr} + \frac{Ze^2}{l+1} - \frac{l+1}{r}\right). \quad (12)$$

The constant $C_3 \equiv G_{nl}$ is determined using recurrence; in fact, from (2) and (12) we have that $\psi_{n1} = A_{n0}^+\psi_{n0} = G_{n0} \times$ function (r) but the normalization condition $\int_0^\infty \psi_{n1}^2 dr = 1$ implies $G_{n0}^2 = n^2/(n^2 - 1)$. Similarly $\psi_{n2} = A_{n1}^+\psi_{n1} = G_{n1}G_{n0} \times$ function (r) together with $\int_0^\infty \psi_{n2}^2 dr = 1$ implies $G_{n1}^2 = n^2/(n^2 - 4)$, and so on. It is not difficult to obtain an expression for C_3 , which is $C_3 = -n[n^2 - (l+1)^2]^{-1/2}$. In this way, equation (12) becomes equivalent to eq. (42) in ref. [11].

In an analogous fashion we can show that the lowering operator

$$A_{nl}^- = \frac{nl}{Ze^2\sqrt{n^2 - l^2}}\left(\frac{d}{dr} - \frac{Ze^2}{l} + \frac{l}{r}\right) \quad (13)$$

satisfies eq. (2).

We have shown that determining the ladder operators A_{nl}^\pm is equivalent to solving a set of linear first-order differential equations. The operators (12) and (13) are equivalent to those obtained by others [10,23] using the factorization method or supersymmetric techniques and, as it is known, they are very useful for evaluating matrix elements of radial functions [24,25].

References

- [1] P R Wallace *Mathematical Analysis of Physical Problems* (New York : Dover) (1984)
- [2] J Morales, J J Peña, M Sánchez and J López-Bonilla *Int J Quantum Chem* **S25** 155 (1991)
- [3] L Infeld and T E Hull *Rev. Mod. Phys.* **23** 21 (1951)
- [4] D S Carlstone *Am. J. Phys* **40** 1459 (1972)
- [5] J N Huffaker and P H Dwivedi *J. Math. Phys.* **16** 862 (1975)
- [6] R Das and A B Sannigrahi *J. Chem. Educ* **58** 383 (1981)
- [7] O L de Lange and R E Raab *Operators Methods in Quantum Mechanics* (Oxford : Clarendon) (1991)
- [8] A Pérez-Lorezana *Rev. Mex. Fís.* **42** 1060 (1996)
- [9] J Douglas and C Peterson *J. Chem. Edu.* **62** 587 (1985)
- [10] Z W Salsburg *Am. J. Phys* **33** 36 (1965)
- [11] J López-Bonilla, D Navarrete, H N Núñez - Yépez and A L Salas-Brito *Int. J. Quantum Chem.* **62** 177 (1997)
- [12] S Y Lee *Am. J. Phys.* **53** 753 (1985)
- [13] R W Haymaker and A R P Rau *Am. J. Phys.* **54** 928 (1986)
- [14] N Aquino, R Gómez, J López-Bonilla, J Morales and D Navarrete *Rev. Mex. Fís* **37** 343 (1991)
- [15] J López-Bonilla, H N Núñez - Yépez and A L Salas-Brito *J Phys.* **B28**, L525 (1995)
- [16] J López-Bonilla, J Morales and M Rosales *Int. J. Quantum Chem* **53** 3 (1995)
- [17] P M Morse *Phys. Rev.* **34** 57 (1929)
- [18] D R Childs *J. Quant. Spectros. Radiat. Transfer* **4** 283 (1964)
- [19] J A Gallas *Phys. Rev.* **A21** 1829 (1980)
- [20] V S Vasan and R J Cross *J. Chem. Phys.* **78** 3869 (1983)
- [21] M Berrondo, A Palma and J López-Bonilla *Int. J. Quantum Chem.* **31** 243 (1987)
- [22] V Gaftoi, J López-Bonilla, J Morales and D Navarrete *Rev Mex. Fís* **36** 649 (1990)
- [23] H R Hauslin *Helv. Phys. Acta* **61** 901 (1988)
- [24] N Bessis and G Bessis *J. Phys.* **A20** 5755 (1987)
- [25] D Yi-Bing *J. Phys.* **A20** 6293 (1987)